

The extreme, workerless inquilines.

***Teleutomyrmex* Kutter, 1950**

- 01) *Teleutomyrmex schneideri* Kutter, 1950
(= *Tetramorium schneideri* (Kutter, 1950), by Ward, Brady, Fisher & Schultz, 2015 ("2014"))
(not *Tetramorium schneideri* Emery, 1898)
(= *Tetramorium inquilinum* Ward, Brady, Fisher & Schultz, 2015 ("2014"))
- 02) *Teleutomyrmex kutteri* Tinaut, 1990
(= *Tetramorium kutteri* (Tinaut, 1990), by Ward, Brady, Fisher & Schultz, 2015 ("2014"))
(not *Tetramorium semilaeve kutteri* Santschi, 1927)
- 03) *Teleutomyrmex seiferti* Kiran & Karaman, in Kiran, et al. 2017
(= *Tetramorium seiferti* (Kiran & Karaman, in Kiran, et al. 2017), by analogy)
- 04) *Teleutomyrmex buschingeri* Lapeva-Gjonova, in Kiran, et al. 2017
(= *Tetramorium buschingeri* (Lapeva-Gjonova, in Kiran, et al. 2017), by analogy)

Not yet described species of extreme, workerless inquiline, from the genus *Teleutomyrmex* Kutter, 1950

- 05) The new, undescribed species from *Teleutomyrmex* Kutter, 1950 from Farab, Turkmenistan... See Dlussky, et al. 1990.

***Anergates* Forel, 1874**

- 06) *Anergates atratulus* (Schenck, 1852), by Forel, 1874
(= *Myrmica atratula* Schenck, 1852)
[Also described as new by Schenck, 1853b]
(= *Tetramorium atratulum* (Schenck, 1852), by Mayr, 1855)
[= *Tomognathus atratulus* (Schenck, 1852), by Mayr, 1863 following Mayr, 1861, obsolete combination.]
(= *Tetramorium atratulum* (Schenck, 1852), by Ward, Brady, Fisher & Schultz, 2015 ("2014"))
- 07) *Anergates friedlandi* Creighton, 1934
[= *Tetramorium friedlandi* (Creighton, 1934), by analogy]

***Tetramorium* Mayr, 1855**

(Only a few species in a big genus.)

08) *Tetramorium microgyna* Santschi, 1918

09) *Tetramorium parasiticum* Bolton, 1980

***Pheidole* Westwood, 1839**

(Only a few species in a big genus.)

10) *Pheidole neokohli* Wilson, 1984

(= *Pheidole kohli* (Wasmann, 1915), by Wilson, 1984)

(= *Anergatides kohli* Wasmann, 1915)

(not *Pheidole kohli* Mayr, 1901)

11) *Pheidole acutidens* (Santschi, 1922), by Wilson, 1984

(= *Bruchomyrma acutidens* Santschi, 1922)

12) *Pheidole argentina* (Bruch, 1932), by Wilson, 1984

(= *Gallardomyrma argentina* Bruch, 1932)

13) *Pheidole parasitica* Wilson, 1984

Excluded from the extreme, workerless inquilines

Once this species was included in the extreme, workerless inquilines but now it is considered to be a workerless inquiline without extreme reductions, e.g. no pupoid males but normal ones. The decision to exclude it was made by Edward Osborne Wilson in 1984 in a study of the inquilines in the genus *Pheidole* Westwood, 1839.

***Pheidole* Westwood, 1839**

(Only one species in a big genus.)

14) *Pheidole kusnezovi* Wilson, 2003

(= *Pheidole symbiotica* (Kusnezov, 1952), by Wilson, 1984)

(= *Eriopheidole symbiotica* Kusnezov, 1952)

(not *Pheidole symbiotica* Wasmann, 1909)

Distribution

01) Europe (Alps, Pyrenees and Northern Spain)

02) Europe (Southern Iberia)

03) Turkey (Anatolia)

04) Europe (Southern Balkans)

05) Turkmenistan

06) Europe

07) North America

08) Southern Africa

09) Southern Africa

10) Central Africa

11) South America

12) South America

13) India

14) South America

Host species

01), 02), 03), 04), 05), 06), 07), 08) and 09) Certain species of the genus *Tetramorium* Mayr, 1855

01) *T. alpestre* Steiner, Schlick-Steiner & Seifert, 2010 and *T. impurum* (Förster, 1850)
and maybe *T. caespitum* (Linnaeus, 1758)?

02) *T. cf. caespitum* (Linnaeus, 1758)

03) *T. cf. chefketi* Forel, 1911

04) *T. cf. chefketi* Forel, 1911

05) A species from the genus *Tetramorium* Mayr, 1855...

06) *T. impurum* (Förster, 1850), *T. caespitum* (Linnaeus, 1758), *T. immigrans* Santschi, 1927
and *T. moravicum* Kratochvil, 1941, *T. diomedium* Emery, 1908, *T. chefketi* Forel, 1911

07) *T. immigrans* Santschi, 1927

08) *T. sericeiventris* Emery, 1877 and *T. sepositum* Santschi, 1918

09) *T. avium* Bolton, 1980

10), 11), 12), 13) and 14) Certain species of the genus *Pheidole* Westwood, 1839

10) *P. megacephala* (Fabricius, 1793) subsp. *melancholica* Santschi, 1912

11) *P. strobili* Emery, 1906

12) *P. nitidula* Emery, 1888

13) *P. indica* Mayr, 1879

14) *P. obscurior* Forel, 1886

A remark about synonymy

Anergates friedlandi Creighton, 1934 is now a synonym from *Anergates atratulus* (Schenck, 1852), more precisely an introduced form in North America. So, the name is *Anergates atratulus* (Schenck, 1852) or, if you follow Ward et al. 2015 ("2014"), *Tetramorium atratulum* (Schenck, 1852)...

Synonyms of host species

- *T. caespitum* (Linnaeus, 1758) (= *Formica caespitum* Linnaeus, 1758)
- *T. impurum* (Förster, 1850) (= *Myrmica impura* Förster, 1850)
- *P. megacephala* (Fabricius, 1793) (= *Formica megecephala* Fabricius, 1793)
- *P. megacephala* (Fabricius, 1793) subsp. *melancholica* Santschi, 1912 was originally described as *P. punctulata* Mayr, 1866 st. *melancholica* Santschi, 1912
- The host species *P. strobili* Emery, 1906 was, in 1922, described as a subspecies of an other species, nl. *P. nitidula* Emery, 1888 subsp. *richteri* Forel, 1909 (a much used junior synonym of the older *P. perversa* Forel, 1908 subsp. *richteri* Forel, 1909) and known as *P. strobili* Emery, 1906 subsp. *richteri* Forel, 1909 since Santschi, 1916. The species *P. strobili* Emery, 1906 was revived from synonymy (with *P. nitidula* Emery, 1888 since Wilson, 2003) and the subspecies was synonymized with it in Casadei-Ferreira et al., 2020.
- *P. obscurior* Forel, 1886 was revived from synonymy with the 1 page older *P. susannae* Forel, 1886 and this in Casadei-Ferreira et al., 2020.